

## Administration Guide

# Intravenous (IV) Push

## Before You Begin


There are important steps you must take before you begin. Please refer to your SASH Teaching Mat for more details.

1. **Clean** your work area.
2. **Wash** your hands.
3. **Gather** your supplies:
  - (2) Pre-filled saline flush syringes
  - (1) Pre-filled heparin flush syringe (if ordered)
  - Medication syringe (at room temperature)
  - Alcohol pads
4. **Inspect** the medication label:
  - Correct patient name
  - Correct medication and dose
  - Medication is not expired
  - Length of infusion
  - The solution should be clear, consistent in color and free of any visible particles

## Prepare Medication

1. Inspect the syringe for any air bubbles. If you see air, remove the bubbles from the syringe by:
  - Removing the tip cap
  - Holding the syringe upright
  - Gently pushing on the plunger until air is removed; try to let only air out of the syringe, and avoid letting medication out.

2. Replace the tip cap on the syringe. Be careful not to touch the inside of the syringe tip cap.




**The tip of the syringe is sterile. Do not** touch it, or let it touch any surfaces. If this happens, discard the syringe as instructed by your nurse and use a new syringe. **Please note** that you will be short one dose and will need more of your medication. Call your pharmacy and explain that you are short one dose of medication. Ask them for directions.

## Flush Your Catheter


Flush your catheter before administering your medication. Please refer to your SASH Teaching Mat for more details.

## Administer Medication

1. Scrub the catheter injection cap with a new alcohol pad for 15 seconds. Let it air dry for five seconds. Be careful not to touch the top of the injection cap with your fingers while you are cleaning it.



2. Remove the cap on top of the syringe. Connect the medication syringe to the catheter injection cap. Then, push the syringe and turn the syringe clockwise until secure.



3. Slowly push in the syringe plunger to administer the medication intravenously over \_\_\_\_\_ minutes. Push \_\_\_\_\_ mL every \_\_\_\_\_ seconds, as instructed by your nurse.
4. Once all the medication has been administered, remove the syringe from the catheter injection cap by twisting counter-clockwise.
5. Discard the medication syringe according to your nurse's instructions.

## Flush Your Catheter

Flush your catheter after administering your medication. Please refer to your SASH Teaching Mat for more details.

### Want to learn more?

Watch a how-to-video on our Resource Center at [coramhc.com](https://coramhc.com).

SASH (SASH=S: Saline; A: Administration of Medication; S: Saline; H: Heparin).

This information is not a substitute for medical advice or treatment. Talk to your doctor or health care provider about this information and any health related questions you have. Coram assumes no liability whatsoever for the information provided or for any diagnosis or treatment made as a result of this information.

Your privacy is important to us. Our employees are trained regarding the appropriate way to handle your private health information.

©2020 Coram LLC. All rights reserved. 75-53178A 112420